



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Factors associated with cytomegalovirus serostatus in young people in England

Citation for published version:

Winter, JR, Taylor, GS, Thomas, OG, Jackson, C, Lewis, JEA & Stagg, HR 2020, 'Factors associated with cytomegalovirus serostatus in young people in England: A cross-sectional study', *BMC Infectious Diseases*, vol. 20, 875. <https://doi.org/10.1186/s12879-020-05572-9>

Digital Object Identifier (DOI):

[10.1186/s12879-020-05572-9](https://doi.org/10.1186/s12879-020-05572-9)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

BMC Infectious Diseases

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Factors associated with cytomegalovirus serostatus in young people in England: A cross-sectional study

Joanne R Winter, PhD,¹ Graham S Taylor, PhD,² Olivia G Thomas, PhD,^{2,3} Charlotte Jackson, PhD,^{1,4} Joanna E A Lewis, PhD,^{5,6} Helen R Stagg PhD^{7*}

1. Centre for Molecular Epidemiology and Translational Research, Institute for Global Health, University College London, London, UK
2. Institute of Immunology and Immunotherapy, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK
3. Current address: Centre for Molecular Medicine, Karolinska University Hospital Solna, 171 76 Stockholm, Sweden
4. Current address: MRC Clinical Trials Unit, University College London, London, UK
5. National Institute for Health Research (NIHR) Health Protection Research Unit in Modelling Methodology and Medical Research Council Centre for Global Infectious Disease Analysis, School of Public Health, Imperial College London, London, UK
6. Current address: Population, Policy and Practice, UCL Great Ormond Street Institute of Child Health, London, UK
7. Usher Institute, University of Edinburgh, Edinburgh, UK

* Corresponding author. Email: helen.stagg@ed.ac.uk

Abstract

Background

Human cytomegalovirus (CMV) is a common herpesvirus which is estimated to infect 83% of the global population. Whilst many infections are asymptomatic, it is an important cause of morbidity and mortality, particularly for immunocompromised people and for infants who are congenitally infected. A vaccine against CMV has been stated as a public health priority, but there are gaps in our understanding of CMV epidemiology. To guide potential future vaccination strategies, our aim was to examine risk factors for CMV seropositivity in young people in England.

Methods

The Health Survey for England (HSE) is an annual, cross-sectional representative survey of households in England during which data are collected through questionnaires, and blood samples are taken. We randomly selected individuals who participated in the HSE 2002, aiming for 25 participants of each sex in each single year age group from 11-24 years. Stored samples were tested for CMV antibodies. We undertook descriptive and regression analyses of CMV seroprevalence and risk factors for infection.

Results

Demographic data and serostatus were available for 732 individuals, of whom 175 (23.7%) were CMV-seropositive. CMV seroprevalence was associated with age, with 18.3% seropositive at 11-14 years compared to 28.3% at 22-24 years. CMV serostatus was also higher in people of non-white ethnicity (adjusted odds ratio [aOR] 6.22, 95% confidence interval [CI] 3.47-11.14), and in adults who were seropositive for EBV (aOR 2.08 [1.06-4.09]). There was no evidence that smoking status, occupation, body mass index and region of England were associated with CMV serostatus.

Conclusions

CMV seroprevalence is strongly associated with ethnicity, and modestly increases with age in 11-24-year-olds. A greater understanding of the transmission dynamics of CMV, and the impact of this on CMV-associated morbidity and mortality, is necessary to inform effective vaccination strategies when a vaccine for CMV becomes available.

50

51 **Key words**

52 Cytomegalovirus, serostatus, transmission, risk factors

Background

The human cytomegalovirus (CMV) is a common human herpesvirus causing lifelong infections, and is estimated to infect 83% of the global population.¹ CMV can be transmitted from symptomatic individuals, via saliva or other body fluids and blood products.² Infection with CMV is typically subclinical in healthy individuals,³ however it is linked to multiple causes of morbidity and mortality. CMV accounts for around 5-8% of infectious mononucleosis cases,^{4,5} and causes disease in immunocompromised people such as transplant and cancer patients.³ Congenitally infected newborns can suffer from cytomegalic inclusion syndrome which can cause long-term neurological damage and, in some cases, life-threatening organ dysfunction.⁵

CMV infection has been associated with shortened life expectancy, particularly in critically ill populations and immunocompromised people (such as those who have undergone organ transplants).⁶ In immunocompetent people in the UK, Gkrania-Klotsas et al. also found that CMV seropositivity was associated with lower life expectancy.⁶ This confirmed the association reported in a population-based cohort study from the US,⁷ as well as populations in older patients⁸ and those with cardiovascular disease.⁹ The association found in Gkrania-Klotsas et al. was specifically with deaths from causes other than cardiovascular disease and cancer, although high levels of CMV IgG antibodies were also associated with cardiovascular mortality.⁶ Other studies in the United States, Finland and United Kingdom found that, in immunocompetent individuals, CMV infection and higher levels of CMV IgG antibodies were linked to higher rates of both cardiovascular⁹ and all-cause mortality,⁷ as well as to cancer incidence¹⁰ and ischemic heart disease.¹¹ However, CMV is negatively associated with multiple sclerosis onset.¹²

In terms of biological pathways, it has been hypothesised that frequent silent reactivations of CMV infection lead to chronic inflammation, which may be a causal factor in the increased risk of mortality.⁶ Additionally, CMV seropositivity has been linked to telomere shortening of T cells, suggesting that CMV may be implicated in immunosenescence, thereby shortening life expectancy.^{2,13} There is also evidence for an immunological phenomenon called 'memory inflation', where a high proportion of CD8+ T cells in older CMV-positive individuals react to an epitope from a CMV protein.¹⁴ This may limit the ability of the immune system to respond to other infections and could be associated with CMV's

ability to infect the vascular endothelium. CMV infection also drives large expansions of cytotoxic virus-specific CD4+ T cells in older individuals, which could ‘take up room’ in the immune system and potentially limit responses to other pathogens.¹⁵

Given the implications of CMV infection, anti-CMV vaccines have been designated high priority by national health agencies, but to date no effective vaccines appear to be imminent.^{16,17} Mathematical modelling of the impact of different vaccination strategies can be used to guide vaccine development efforts and will be necessary to inform the optimal strategies for deployment of such a vaccine if or when it becomes available. A thorough understanding of CMV epidemiology is necessary for the development of such models.

CMV seroprevalence increases with age, and infection occurs at younger ages in economically developing countries,^{2,6} possibly due to higher rates of breastfeeding than in the UK (CMV is known to be transmitted through breast milk). A large population-based UK cohort study found that CMV infection was more common in women than in men.⁶ Lower income and education levels, and ethnicities other than white, have been associated with earlier age at CMV infection.¹⁸ CMV infection is also correlated with EBV infection.^{19,20} Socioeconomic status is strongly correlated with CMV infection; the reasons for this could include larger family size,²¹ or have been hypothesised to be a result of stress induced by low socioeconomic status contributing to the down-regulation of the immune system and increased susceptibility to infection.¹⁸

To date in the UK, studies of CMV seroprevalence have focused on older adults,⁶ pregnant women,²² and young children.^{23,24} However transmission can occur at all ages, and the association of CMV with infectious mononucleosis suggests that infection during adolescence could also be an important cause of morbidity. Therefore, our aim was to investigate the sociodemographic and lifestyle factors, particularly age, associated with CMV serostatus in children and young adults in England, in order to gain a better understanding of the epidemiology of CMV in this age group.

Methods

Study population

The Health Survey for England (HSE) is a cross-sectional annual representative survey of English households. The methods have been described previously.²⁵ As part of a larger study

investigating Epstein-Barr virus infection and transmission,^{20,26} we used data from randomly selected participants in the 2002 HSE, aiming for 25 male and 25 female participants in each single-year age group from 11-24 years.

Outcome: Seropositivity for Cytomegalovirus infection

We used stored blood serum samples collected by the HSE. Commercial ELISA kits from EUROIMMUN, Germany (EI2791-9601-G, EI2570-9601G) were used to detect CMV-specific IgG and EBV virus capsid antigen (VCA)-specific IgG from the serum samples. Assays were conducted according to the manufacturer's instructions, and we calculated serum antibody concentrations using a standard curve. Results were presented in relative units (RU/mL) using the following thresholds; samples of <16RU/mL were classed as negative, ≤16 to <22RU/mL were borderline, and ≥22RU/mL were positive. For the analyses presented here, borderline results (CMV n=1, EBV n=5) were considered seropositive.²⁰

Statistical Analysis

We used Stata version 15.0 for data analysis. Stata's svy commands were used to weight our sample to be representative of the age and sex of the 2002 English population, using data from the Office for National Statistics.²⁷ All stated percentages are weighted. We conducted descriptive analyses of the study population. We used ArcMap 10.3.1 to map CMV seroprevalence by English Government Office Region.²⁸

We used logistic regression models to investigate factors associated with being seropositive for CMV. We used a causal inference framework to identify *a priori* factors that needed to be included in multivariable models, drawing on the available data from the HSE. This resulted in two multivariable regression models. We built a 'whole-population' model, which included our entire study population, to examine the following factors: age, sex, ethnicity (categorised as 'white' or 'other' due to small numbers of non-white participants), body mass index (BMI; categorised as 'underweight' [BMI <20], 'healthy weight' [20-25], 'overweight' [25-30] or 'obese' [>30]), region of England and EBV serostatus. Additionally, we built a second 'adults-only' model, which only included participants aged ≥16 years, and additionally included data from questions which were only asked to adults; smoking status (never smoked, current smoker, smoked in past) and occupational category (higher managerial and professional, intermediate occupations, routine and manual occupations,

never worked or long-term unemployed, and other). Individuals missing data on one or more variables were excluded from the regression modelling.

Ethical approval

This study was approved by the University College London Research Ethics Committee (5683/002). The HSE obtained informed written consent from participants at the time of recruitment for blood samples to be collected and stored for future analyses.[15] A parent/guardian of participants also provided written consent for the interviewing of participants who were younger than 16 years, and for the taking of blood samples from participants who were younger than 18 years.

Results

Our study sample included 732 individuals aged 11-24 years, of whom 175 (23.7%) were CMV-seropositive. Seroprevalence by participant characteristics are shown in Table 1. There was a slight increase in CMV seropositivity associated with age, from 18.3% at 11-14 years to 28.3% at 22-24 years. CMV seroprevalence was much lower in white people (19.2%) than people of other ethnicities (61.6%). Considerable variation in CMV seroprevalence was observed by region of England (Figure 1, Table 1), being highest in London (47.9%) and otherwise varying between 16.7% in the south-east and 28.7% in the east of England. CMV serostatus was also higher in women (26.9%) than in men (20.4%) and people who were EBV-seropositive (25.7%) than EBV-seronegative (17.9%).

Univariable and multivariable regression models were built examining the factors associated with CMV positivity. Factors associated with CMV seropositivity were largely consistent between the univariable and multivariable models (Table 2), although confidence intervals tended to be wider in the multivariable models. Ethnicities other than white were strongly associated with CMV seropositivity in both univariable and multivariable models (adjusted odds ratio [aOR] 6.22, 95% confidence interval [CI] 3.47-11.14). EBV serostatus was associated with CMV serostatus in the univariable model (odds ratio [OR] 1.59 [1.06-2.39] and in the adults-only multivariable model (aOR 2.08 [1.06-4.09]), but not in the multivariable model which included children (aOR 1.21 [0.76-1.92]). Female sex was associated with higher CMV positivity in the univariable model (OR 1.44, 1.02-2.02); the multivariable models had similar point estimates, but the confidence intervals included

unity. Region of England was not associated with CMV serostatus in multivariable models. Neither smoking status nor occupation were associated with CMV serostatus in adults.

Our study sample included 732 individuals aged 11-24 years, of whom 175 (23.7%) were CMV-seropositive. The characteristics of seropositive individuals are shown in Table 1. Univariable and multivariable regression models were built examining the factors associated with CMV positivity. Factors associated with CMV seropositivity were largely consistent between the univariable and multivariable models (Table 2), although confidence intervals tended to be wider in the multivariable models.

There was an increase in CMV seropositivity associated with age, from 18.3% at 11-14 years to 28.3% at 22-24 years, but the confidence intervals between strata overlapped in logistic regression models. CMV seroprevalence was much higher in people of non-white ethnicities than in white people (61.6% vs 19.2%; aOR 6.22, 95% CI 3.47-11.14). CMV serostatus was also higher in women (26.9%) than in men (20.4%); the CI for this association excluded unity in a univariable model (odds ratio [OR] 1.44, 95% confidence interval [CI] 1.02-2.02); the multivariable models had similar point estimates, but the confidence intervals included unity. Neither smoking status nor occupation were associated with CMV serostatus in adults.

CMV seropositivity was higher in people who were EBV-seropositive (25.7%) than EBV-seronegative (17.9%). EBV serostatus was associated with CMV serostatus in the univariable model (OR 1.59 [1.06-2.39] and in the adults-only model (aOR 2.08 [1.06-4.09]), but not in the multivariable model which included children (aOR 1.21 [0.76-1.92]).

Considerable variation in CMV seroprevalence was observed by region of England (Figure 1, Table 1), CMV seroprevalence was highest in London (47.9%) and otherwise varied between 16.7% in the south-east to 28.7% in the east of England. However, region of England was not associated with CMV serostatus in multivariable models.

Discussion

In this study of young people in England, we found that just under a quarter of people aged 11-24 years were infected with CMV, and that seroprevalence increased over this age range. CMV infection was also strongly correlated with non-white ethnicity and more weakly

associated with EBV infection. There was no association observed between CMV and region of England, smoking status, BMI, or occupation.

CMV and EBV serostatus were positively associated in univariable analyses, and when the multivariable analysis was restricted to adults, but not in the multivariable model which also included children aged 11-15 years. As discussed in our previous paper,²⁰ both CMV and EBV are associated with increasing age, however EBV increases more rapidly during adolescence than CMV. Thus, in a whole-cohort model adjusting for age, this association may not be visible. As the associations between age and both CMV and EBV²⁰ are less strong in adults, it is possible that there was enough of a residual effect that the association between CMV and EBV could be detected in the adults-only model.

Given the cross-sectional nature of our study, the relative temporality of the two infections could not be assessed. Although established, the relationship between CMV and EBV is not well understood. It is known that EBV seroprevalence is higher than CMV seroprevalence in all age groups and that both increase with age,²⁰ but it is not known whether this relationship is causal or whether the association results from shared genetic, immunological and/or sociodemographic risk factors. Longitudinal studies with serial testing and a larger sample size would be necessary to explore this association in more detail.

We observed a strong association between ethnicity and CMV seroprevalence; the odds of being CMV positive were approximately seven times higher for people of ethnicities other than white than the odds for white people. This may be the result of different social mixing patterns, larger households, different eating or hygiene habits, lower breastfeeding rates in white people (resulting in less vertical transmission of CMV through breastmilk), possibly different countries of birth (of participants or their parents) or residual confounding of socioeconomic status. This strong association with ethnicity is also likely to be a confounder in the association between CMV and region, particularly London, that was only observed in univariable models, as there is a higher proportion of ethnic minorities living in London than elsewhere in England.²⁹ We were unable to analyse associations with ethnicity in more detail due to small numbers of participants; the “non-white” group comprised 57% Asian/Asian British (n=44), 19% black/black British (n=15), 14% mixed ethnicity (n=11) and 9% other ethnicity (n=7). Further study of the association of ethnicity with CMV seroprevalence is needed in diverse cohorts.

Our study benefits from a sample drawn from a highly rigorous, annual, representative survey of people in England, which we weighted to be representative of the English population, and the use of a quality-managed commercial assay to measure the antibody response. The limitations of our work include the use of a cross-sectional study design, preventing determination of the temporality of certain associations, and the age of the data; 2002 was the most recent year for which the HSE collected consent to analyse blood samples for blood-borne viruses. More recent data from the UK biobank found that 58% of those aged 40-69 years were seropositive for CMV at enrolment (2006-2010),³⁰ and as CMV seroprevalence increases with age throughout life, the prevalence observed in young people in our study is consistent with what could be expected. An older study examined CMV seroprevalence in 1991 and 2002 and found that prevalence in young people did not differ between these two timepoints,³¹ and so there is no particular reason to believe CMV seroprevalence has changed substantially since then. We also consider it unlikely that the associations between CMV and the risk factors we studied would have changed substantially since 2002, and therefore the associations we observed are likely to be consistent today even if there had been a slight change in CMV seroprevalence.

The relatively low seroprevalence of CMV meant this study may have lacked power to detect associations, particularly in the multivariable models. We were also limited in the variables that were available, we were unfortunately unable to examine associations with household size or household income. Additionally, the geographical variables available were lacking in granularity, meaning we were not able to explore regional differences in more depth or examine whether regional variation was associated with other sociodemographic risk factors.

We observed only a modest increase in CMV seroprevalence associated with age, suggesting that adolescence is not a key transmission period for CMV as it is for EBV (for which seroprevalence increases from 60% in 11-14 year olds to 93% in 22-24 year olds²⁰). This may contribute to the lower incidence of CMV-associated (versus EBV-associated) infectious mononucleosis.^{4,5} Previous studies have shown that 15% of white British and 44% of British Pakistani infants were infected with CMV by the age of 2 years, and that seroprevalence was 59% in an adult cohort aged 40-79 years. In combination with our results, this suggests that after early childhood, there is no 'key' age group in which CMV seroprevalence sharply

increases, and that infection continues to increase during adulthood, particularly for white British individuals. A better understanding of the interactions between age at CMV infection, and the development of CMV-related morbidity and mortality, is necessary to be able to develop an appropriate vaccination strategy, when a vaccine becomes available.

Conclusions

CMV seroprevalence is strongly associated with ethnicity, and modestly increases with age. A greater understanding of the transmission dynamics of CMV, and the impact of this on CMV-associated morbidity and mortality, is necessary to inform effective vaccination strategies when a vaccine for CMV becomes available.

Declarations

Ethics approval and consent to participate

This study was approved by the University College London Research Ethics Committee (5683/002). The HSE obtained informed written consent for blood samples to be collected and stored for future analyses.

List of abbreviations

aOR: adjusted odds ratio

BMI: body mass index

CI: confidence interval

CMV: cytomegalovirus

EBV: Epstein-Barr virus

HSE: health survey for England

OR: odds ratio

RU: relative units

UK: United Kingdom

VCA: virus capsid antigen

Consent for publication

Not applicable

Availability of data and materials

The data used in this study was under license from the Health Survey for England, and so are not publicly available, but can be requested from the HSE.

Competing interests

GT reports personal fees from Genocea Biosciences, outside the submitted work. CJ is an Associate Editor at BMC Public Health. All other authors have no competing interests.

Funding

This work was supported by the Wellcome Trust [204419]. The funding source has no role in the study design, collection, analysis or interpretation of the data, the writing of the paper or the decision to submit for publication. The corresponding author had full access to all data in the study and had final responsibility to submit the paper for publication.

Authors' contributions

HRS, JL and GT designed the study. OT and GT conducted the serological testing. JRW conducted the data analysis and drafted the paper. JRW, CJ, HRS, JL and GT interpreted the results. All authors critically revised the paper and approved the final version for publication.

Acknowledgements

We thank our colleagues at UCL and NatCen Social Research, and the interviewers, research nurses and participants of the Health Survey for England, and Shaun Scholes for assistance weighting the HSE data for analysis.

References

1. Zuhair, M. *et al.* Estimation of the worldwide seroprevalence of cytomegalovirus: A systematic review and meta-analysis. *Rev. Med. Virol.* **29**, e2034 (2019).
2. Arens, R., Remmerswaal, E. B. M., Bosch, J. A. & Lier, R. A. W. van. 5th International Workshop on CMV and Immunosenescence - A shadow of cytomegalovirus infection on immunological memory. *Eur. J. Immunol.* **45**, 954–957 (2015).
3. Navarro, D. Expanding role of cytomegalovirus as a human pathogen: Cytomegalovirus and Human Disease. *J. Med. Virol.* **88**, 1103–1112 (2016).

- 320 4. Evans, A. Infectious mononucleosis and related syndromes. *Am. J. Med. Sci.* **276**, 325–
321 340 (1978).
- 322 5. Landolfo, S., Gariglio, M., Gribaudo, G. & Lembo, D. The human cytomegalovirus.
323 *Pharmacol. Ther.* **98**, 269–297 (2003).
- 324 6. Gkrania-Klotsas, E. *et al.* Seropositivity and Higher Immunoglobulin G Antibody Levels
325 Against Cytomegalovirus Are Associated With Mortality in the Population-Based
326 European Prospective Investigation of Cancer–Norfolk Cohort. *Clin. Infect. Dis.* **56**, 1421–
327 1427 (2013).
- 328 7. Simanek, A. M. *et al.* Seropositivity to cytomegalovirus, inflammation, all-cause and
329 cardiovascular disease-related mortality in the United States. *PloS One* **6**, e16103 (2011).
- 330 8. Wang, G. C. *et al.* Cytomegalovirus infection and the risk of mortality and frailty in older
331 women: a prospective observational cohort study. *Am. J. Epidemiol.* **171**, 1144–1152
332 (2010).
- 333 9. Strandberg, T. E., Pitkala, K. H. & Tilvis, R. S. Cytomegalovirus antibody level and
334 mortality among community-dwelling older adults with stable cardiovascular disease.
335 *JAMA* **301**, 380–382 (2009).
- 336 10. Lepiller, Q., Tripathy, M. K., Di Martino, V., Kantelip, B. & Herbein, G. Increased HCMV
337 seroprevalence in patients with hepatocellular carcinoma. *Viol. J.* **8**, 485 (2011).
- 338 11. Gkrania-Klotsas, E. *et al.* Higher Immunoglobulin G Antibody Levels Against
339 Cytomegalovirus Are Associated With Incident Ischemic Heart Disease in the Population-
340 Based EPIC-Norfolk Cohort. *J. Infect. Dis.* **206**, 1897–1903 (2012).
- 341 12. Sundqvist, E. *et al.* Cytomegalovirus seropositivity is negatively associated with multiple
342 sclerosis. *Mult. Scler. Houndmills Basingstoke Engl.* **20**, 165–173 (2014).
- 343 13. van de Berg, P. J. E. J. *et al.* Cytomegalovirus infection reduces telomere length of the
344 circulating T cell pool. *J. Immunol. Baltim. Md 1950* **184**, 3417–3423 (2010).
- 345 14. Hosie, L. *et al.* Cytomegalovirus-Specific T Cells Restricted by HLA-Cw*0702 Increase
346 Markedly with Age and Dominate the CD8+ T-Cell Repertoire in Older People. *Front.*
347 *Immunol.* **8**, (2017).
- 348 15. Pachnio, A. *et al.* Cytomegalovirus Infection Leads to Development of High Frequencies
349 of Cytotoxic Virus-Specific CD4+ T Cells Targeted to Vascular Endothelium. *PLoS Pathog.*
350 **12**, e1005832 (2016).

16. Sung, H. & Schleiss, M. R. Update on the current status of cytomegalovirus vaccines. *Expert Rev. Vaccines* **9**, 1303–1314 (2010).
17. Modlin, J. F. *et al.* Vaccine Development to Prevent Cytomegalovirus Disease: Report from the National Vaccine Advisory Committee. *Clin. Infect. Dis.* **39**, 233–239 (2004).
18. Dowd, J. B., Aiello, A. E. & Alley, D. E. Socioeconomic Disparities in the Seroprevalence of Cytomegalovirus Infection in the U.S. Population: NHANES III. *Epidemiol. Infect.* **137**, 58–65 (2009).
19. Levine, H. *et al.* Seroepidemiology of Epstein-Barr virus and cytomegalovirus among Israeli male young adults. *Ann. Epidemiol.* **22**, 783–788 (2012).
20. Winter, J. R. *et al.* Predictors of Epstein-Barr virus serostatus in young people in England. *BMC Infect. Dis.* **19**, 1007 (2019).
21. Lachmann, R. *et al.* Cytomegalovirus (CMV) seroprevalence in the adult population of Germany. *PLoS ONE* **13**, (2018).
22. Pembrey, L. *et al.* Seroprevalence of Cytomegalovirus, Epstein Barr Virus and Varicella Zoster Virus among Pregnant Women in Bradford: A Cohort Study. *PLoS ONE* **8**, (2013).
23. Pembrey, L. *et al.* Cytomegalovirus, Epstein-Barr virus and varicella zoster virus infection in the first two years of life: a cohort study in Bradford, UK. *BMC Infect. Dis.* **17**, 220 (2017).
24. Pembrey, L., Waiblinger, D., Griffiths, P. & Wright, J. Age at cytomegalovirus, Epstein Barr virus and varicella zoster virus infection and risk of atopy: The Born in Bradford cohort, UK. *Pediatr. Allergy Immunol.* **30**, 604–613 (2019).
25. Deverill, C. *et al.* *Health Survey for England 2002: The Health of Children and Young People. Methodology & Documentation.* (The Stationary Office, 2002).
26. Goscé, L., Winter, J. R., Taylor, G. S., Lewis, J. E. A. & Stagg, H. R. Modelling the dynamics of EBV transmission to inform a vaccine target product profile and future vaccination strategy. *Sci. Rep.* **9**, 1–9 (2019).
27. Office for National Statistics. Estimates of the population for the UK, England and Wales, Scotland and Northern Ireland. <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/datasets/populationestimatesforukenglandandwalesscotlandandnorthernireland> (2018).

382 28. Office for National Statistics. Census boundary data [United Kingdom].
383 <https://census.ukdataservice.ac.uk/get-data/boundary-data.aspx> (2011).
384 29. UK Data Service. Census aggregate data. [https://census.ukdataservice.ac.uk/get-](https://census.ukdataservice.ac.uk/get-data/aggregate-data.aspx)
385 [data/aggregate-data.aspx](https://census.ukdataservice.ac.uk/get-data/aggregate-data.aspx) (2001).
386 30. UKB : Data-Field 23054. <https://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=23054>.
387 31. Vyse, A. J., Hesketh, L. M. & Pebody, R. G. The burden of infection with cytomegalovirus
388 in England and Wales: how many women are infected in pregnancy? *Epidemiol. Infect.*
389 **137**, 526–533 (2009).
390
391
392

393 **Tables**394 **Table 1: The baseline characteristics of the study population and number and weighted**
395 **percentage of individuals seropositive for CMV in England in 2002.**

Variable	Total number	Number CMV seropositive (weighted %)
Total	732	175 (23.7)
Sex		
Male	364	74 (20.4)
Female	368	101 (26.9)
Age at last birthday (years)		
11-14	208	39 (18.3)
15-18	212	52 (24.6)
19-21	156	40 (25.6)
22-24	156	44 (28.3)
Ethnicity		
White	665	127 (19.2)
Other	77	48 (61.6)
BMI		
Underweight	60	17 (28.1)
Healthy weight	418	94 (22.0)
Overweight	141	34 (23.8)
Obese	87	25 (29.9)
Missing	26	5 (19.4)
EBV serostatus		
EBV-seronegative	547	34 (17.9)
EBV-seropositive	185	141 (25.7)

Region of England

East of England	78	24 (28.7)
North East	34	6 (18.0)
North West	130	27 (19.7)
Yorkshire and The Humber	82	18 (22.1)
East Midlands	74	13 (17.0)
West Midlands	70	16 (23.4)
London	63	30 (47.9)
South East	119	20 (16.7)
South West	82	21 (26.4)

Smoking status*

Never smoked	86	52 (27.9)
Current smoker	134	35 (26.0)
Past smoker	147	36 (24.5)
Missing	5	3 (60.0)

Occupational category*

Higher managerial and professional	83	26 (31.6)
Intermediate occupations	69	20 (28.9)
Routine and manual occupations	254	61 (24.0)
Never worked or long-term unemployed	11	2 (18.1)
Other	55	17 (30.3)

*Adults aged ≥ 16 years only (n=472). Percentages account for the weighting of the sample to be representative of the English population in 2002 with respect to age and sex. BMI: body mass index, CI: confidence interval, CMV: cytomegalovirus, EBV: Epstein-Barr virus.

Table 2: Univariable and multivariable logistic regression models of factors associated with Cytomegalovirus seropositivity in England in 2002.

	Whole-population		Adults only*
	Univariable OR (95% CI)	Multivariable aOR (95% CI)	Multivariable aOR (95% CI)
Sex			
Male	1.00	1.00	1.00
Female	1.44 (1.02-2.02)	1.39 (0.95-2.05)	1.50 (0.95-2.35)
Age at last birthday (years)			
11-14	1.00	1.00	
15-18	1.46 (0.89-2.40)	1.55 (0.88-2.75)	1.00
19-21	1.54 (0.89-2.66)	2.24 (1.21-4.14)	1.05 (0.58-1.91)
22-24	1.76 (1.04-2.99)	1.77 (0.95-3.29)	0.74 (0.38-1.46)
Ethnicity			
White	1.00	1.00	1.00
Other	6.75 (4.23-10.77)	6.22 (3.47-11.14)	6.98 (3.18-15.32)
BMI			

Underweight	1.39 (0.77-2.52)	1.18 (0.62-2.24)	0.98 (0.50-1.93)
Healthy weight	1.00	1.00	1.00
Overweight	1.11 (0.71-1.74)	1.15 (0.71-1.86)	0.84 (0.47-1.53)
Obese	1.52 (0.90-2.56)	1.50 (0.81-2.77)	1.06 (0.40-2.76)
EBV serostatus			
Negative	1.00	1.00	1.00
Positive	1.59 (1.06-2.39)	1.21 (0.76-1.92)	2.08 (1.06-4.09)
Region of England			
East of England	1.00	1.00	1.00
North East	0.54 (0.18-1.66)	0.51 (0.17-1.50)	0.57 (0.16-2.02)
North West	0.61 (0.31-1.19)	0.62 (0.31-1.22)	0.60 (0.25-1.40)
Yorkshire and The Humber	0.70 (0.33-1.51)	0.67 (0.30-1.51)	0.62 (0.26-1.47)
East Midlands	0.51 (0.21-1.19)	0.52 (0.21-1.26)	0.70 (0.24-2.04)
West Midlands	0.76 (0.35-1.63)	0.81 (0.36-1.79)	0.53 (0.18-1.60)
London	2.28 (1.13-4.61)	1.18 (0.52-2.69)	1.28 (0.45-3.61)
South East	0.50 (0.24-1.03)	0.43 (0.20-0.93)	0.44 (0.17-1.14)

South West	0.89 (0.46-1.70)	1.09 (0.55-2.15)	0.72 (0.29-1.78)
------------	------------------	------------------	------------------

Smoking status

Never smoked	1.00	-	1.00
Current smoker	0.91 (0.54-1.51)	-	1.08 (0.58-2.01)
Smoked in past	0.84 (0.51-1.38)	-	0.96 (0.54-1.69)


Occupational category

Higher managerial and professional	1.00	-	1.00
Intermediate occupations	0.88 (0.44-1.76)	-	0.66 (0.31-1.42)
Routine and manual occupations	0.68 (0.40-1.18)	-	0.63 (0.32-1.24)
Never worked or long-term unemployed	0.48 (0.09-2.45)	-	0.40 (0.07-2.14)
Other	0.94 (0.46-1.93)		0.47 (0.17-1.28)

*Adults aged ≥16 years only (n=472). †16-18 years for 'adult-only' model. Odds ratios account for the weighting of the sample to be representative of the English population in 2002 with respect to age and sex. The 'whole population' multivariable model included age, sex, CMV serostatus, ethnicity, BMI and region of England. The 'adults only' multivariable model included all variables shown in the table. aOR: adjusted odds ratio, BMI: body mass index, CI: confidence interval, CMV: cytomegalovirus, OR: unadjusted odds ratio.

Figures

Figure 1: Weighted Cytomegalovirus seroprevalence by English Government Office Region in 2002.



Contains National Statistics data © Crown copyright and database right [2011] Contains public sector 6 information licensed under the Open Government Licence v3.0